

Death rate due to STDs in the US, 1996 - 2014


ARGON

DATA VISUALIZATION

(C) 2019 Novixys Software, Inc.

Death rate due to STDs in the US, 1996 - 2014

Deaths per 100,000 people due to sexually-transmitted diseases by age group.



Deaths due to Sexually Transmitted Diseases in the US, 1996 - 2014

Chlamydia was the most common cause of death due to STDs in the US. Overall, between 1996 and 2014 the death rate was 341 per 100,000, followed by 115 for Gonorrhea and 4 deaths for Syphilis. The death rate for 20-24 year olds is especially high at 1800 deaths for Chlamydia, 524 for Gonorrhea and 9 deaths for Syphilis.

[See this page for an annual breakdown](#) of the numbers for Chlamydia.

Input Data

Disease	Age	RatePer100k
Syphilis	0-14 years	0.04
Gonorrhea	0-14 years	8.33
Chlamydia	0-14 years	25.74
Syphilis	15-19 years	3.31
Gonorrhea	15-19 years	442.39
Chlamydia	15-19 years	1636.71
Syphilis	20-24 years	9.27
Gonorrhea	20-24 years	523.50
Chlamydia	20-24 years	1799.41
Syphilis	25-29 years	8.97
Gonorrhea	25-29 years	274.96
Chlamydia	25-29 years	743.25
Syphilis	30-34 years	7.68
Gonorrhea	30-34 years	148.60
Chlamydia	30-34 years	309.05
Syphilis	35-39 years	7.07
Gonorrhea	35-39 years	90.11
Chlamydia	35-39 years	139.89
Syphilis	40+ years	2.51
Gonorrhea	40+ years	20.50
Chlamydia	40+ years	22.88
Syphilis	Unknown	0.00
Gonorrhea	Unknown	0.00
Chlamydia	Unknown	0.00

Sources

- CDC Wonder, <https://wonder.cdc.gov/>